

Performance and Training Standards for Endovascular Ischemic Stroke Treatment

Philip M. Meyers, MD, H. Christian Schumacher, MD, Michael J. Alexander, MD, Colin P. Derdeyn, MD, Anthony J. Furlan, MD, Randall T. Higashida, MD, Christopher J. Moran, MD, Robert W. Tarr, MD, Donald V. Heck, MD, Joshua A. Hirsch, MD, Mary E. Jensen, MD, Italo Linfante, MD, Cameron G. McDougall, MD, Gary M. Nesbit, MD, Peter A. Rasmussen, MD, Thomas A. Tomsick, MD, Lawrence R. Wechsler, MD, John R. Wilson, MD, and Osama O. Zaidat, MD, MS

Stroke is the third-leading cause of death in the United States, Canada, Europe, and Japan. According to the American Heart Association and the American Stroke Association, 750,000 new strokes occur each year, resulting in 200,000 deaths (or 1 of every 16 deaths) per year in the United States alone. Endovascular therapy for patients with acute ischemic stroke is an area of intense investigation. The American Stroke Association has given a qualified endorsement of intra-arterial (IA) thrombolysis in selected patients. IA thrombolysis has been studied in 2 randomized trials and numerous case series. Although 2 devices have been granted FDA 3 approval with an indication for mechanical stroke thrombectomy, none of these devices has demonstrated efficacy in improving patient outcomes. This report defines what constitutes adequate training to perform neuroendovascular procedures in patients with acute ischemic stroke and identifies the performance standards that should be adopted to assess outcomes. These guidelines have been written and approved by multiple neuroscience societies that historically have been directly involved in the medical, surgical, and endovascular care of patients with acute stroke, including the Neurovascular Coalition and its participating societies: the Society of NeuroInterventional Surgery; American Academy of Neurology; American Association of Neurological Surgeons, Cerebrovascular Section; and Society of Vascular & Interventional Neurology. **Key Words:** Stroke—endovascular therapy—training standards—performance—intra-arterial—fellowship—endovascular.

© 2009 by American Academy of Neurology, American Association of Neurological Surgeons Cerebrovascular Section, Society of NeuroInterventional Surgery, and Society of Vascular & Interventional Neurology

From the Writing Group for the American Academy of Neurology, American Association of Neurological Surgeons Cerebrovascular Section, Society of NeuroInterventional Surgery, and Society of Vascular & Interventional Neurology.

Received June 16, 2009; revision received August 12, 2009; accepted August 15, 2009.

The four societies hold the full copyright for the position statement as depicted in this manuscript.

Address correspondence to Philip M. Meyers, Associate Professor, Radiology and Neurological Surgery, New York Presbyterian

Hospital, Columbia University, College of Physicians & Surgeons, Neurological Institute, 710 West 168th Street, Suite 428, New York, NY 10032. E-mail: pmm2002@columbia.edu.

1052-3057/\$—see front matter

© 2009 American Academy of Neurology, American Association of Neurological Surgeons Cerebrovascular Section, Society of NeuroInterventional Surgery, and Society of Vascular & Interventional Neurology.

doi:10.1016/j.jstrokecerebrovasdis.2009.08.001

Stroke is the third-leading cause of death in the United States, Canada, Europe, and Japan. According to the American Heart Association and the American Stroke Association, 750,000 new strokes occur each year, resulting in 200,000 deaths (or 1 of every 16 deaths) per year in the United States alone.¹ Ischemic stroke accounts more than 80% of the total number of strokes, with hemorrhagic stroke comprising the remainder. Stroke is the leading cause of adult disability in North America² and the primary cause for in-patient Medicare reimbursement for long-term adult care.^{3,4} The National Institutes of Health estimates that the cost of stroke now exceeds \$62 billion annually.¹

At present, the only therapy demonstrated to improve clinical outcomes from acute ischemic stroke is thrombolysis of the clot responsible for the ischemic event.⁵ Specifically, the only FDA-approved therapy for stroke is intravenous tissue plasminogen activator (tPA) administered within 3 hours of stroke onset.⁵ Endovascular therapy for patients with acute ischemic stroke is an area of intense investigation. The American Stroke Association has given a qualified endorsement of intra-arterial (IA) thrombolysis in selected patients. IA thrombolysis has been studied in 2 randomized trials^{6,7} and numerous case series that have demonstrated clinical efficacy for IA prourokinase in proximal middle cerebral artery occlusion.⁷ Although 2 devices have been granted FDA approval with an indication for mechanical stroke thrombectomy,⁸ neither of these devices has demonstrated efficacy in improving patient outcomes despite a high rate of recanalization awaiting randomized trials.

In this report we define what constitutes adequate training to perform neuroendovascular procedures in patients with acute ischemic stroke and identify the performance standards that should be adopted to assess outcomes. Neuroendovascular procedures are technically challenging and not directly transferable from other vascular systems, involve an organ with unique physiology and anatomy, and require careful patient selection because of the risk of potentially fatal brain hemorrhage. Studies of emerging technologies over the past 20 years have made it abundantly clear that inadequate physician training and experience can adversely affect clinical outcomes.^{9,10} Overexuberance by both physicians and industry has led unqualified physicians to perform endovascular carotid revascularization procedures, yielding inferior results.^{9,10} Thus, it is especially important that the involved physicians strictly adhere to appropriate standards when performing high-risk procedures such as those involved in treating endovascular stroke. Standardization of training requirements is critically important as the interest in and utilization of endovascular methods increases among various specialties.

These guidelines are modeled after earlier standards documents, such as the training, competency, and credentialing standards for diagnostic cerebral angiography, carotid stenting, and treatment of acute stroke, written and

endorsed by multispecialty groups¹¹⁻¹³ and training standards for the performance of uterine artery embolization written by the Society of Interventional Radiology.¹⁴ These guidelines also parallel the training standards of successful subspecialty training programs such as interventional cardiology, and the credentialing standards for the performance of acute coronary interventions.¹⁵ These guidelines have been written and approved by multiple neuroscience societies that historically have been directly involved in the medical, surgical, and endovascular care of patients with acute stroke and considered expert in the field of endovascular stroke therapy, including the Society of NeuroInterventional Surgery; American Academy of Neurology; American Association of Neurological Surgeons, Cerebrovascular Section; and Society of Vascular & Interventional Neurology.

Minimum Training Requirement for Acute Stroke Interventions

Cognitive training and qualifications:

1. Accreditation Council for Graduate Medical Education (ACGME)-approved residency training including documented cerebrovascular training, including the diagnosis and management of acute stroke and the interpretation of cerebral arteriography and brain imaging under the supervision of a board-certified neurologist, neurosurgeon, or neuroradiologist with the American Board of Medical Speciality (ABMS) eligibility or certification. A minimum of 6 months during a 4-year residency is suggested.
2. One year of graduate medical education in endovascular surgical neuroradiology. An ACGME-approved program is preferred but not required.

Technical training and qualifications:

1. Documented prior training and experience in catheter arteriography, including 100 cerebral arteriograms. Clinical outcomes must meet or exceed the American College of Radiology (ACR) benchmarks for technical success and complications.^{11,16}
2. Documented prior training and experience in intracranial microcatheter (≤ 3 French) and microguidewire (≤ 0.014 inch) navigation under the supervision of fellowship-trained and credentialed neurointerventionalist(s).
3. Documented prior experience in assessment and performance of endovascular stroke interventional procedures as the primary operator in 10 patients under the supervision of fellowship-trained and credentialed neurointerventionalists(s).
4. Previously credentialed physicians who perform IA catheter-directed stroke procedures at their local institutions should have documented procedural

and clinical outcomes that meet national standards and published evidence-based guidelines.^{12,13}

During the next 2–3 years, as the nation progresses to regional stroke care with comprehensive stroke centers, it is conceivable that selective hospitals may choose to credential individuals for IA stroke therapy who have not had a full year of neurointerventional fellowship training. While this is not endorsed as best clinical practice, if such local situations occur and if hospitals choose to credential individuals who have not had a full year of neurointerventional training, then it is recommended that at a minimum these individuals should receive adequate neuroscience cognitive training, including a minimum of 6 months documented neuroscience training in an ACGME-approved postgraduate training program; adequate technical and interpretive cerebral angiography training, including a minimum primary operator experience of 100 appropriately supervised cerebral angiograms, which is the minimum prerequisite for neurointerventional training; adequate cerebral microcatheter experience, including a minimum of 30 cases as primary operator in the intracranial internal carotid artery/vertebral basilar circulation, mentored by a credentialed fellowship-trained neurointerventionalist; and mentored experience in IA stroke therapy, including a minimum of 10 cases mentored by a credentialed fellowship-trained neurointerventionalist credentialed in IA stroke therapy, before being credentialed to perform IA therapy.

It is anticipated that by 2012, stroke centers that are providing IA stroke therapy will be staffed exclusively with fellowship-trained neurointerventionalist and/or practitioners who have met the foregoing prerequisite guidelines and have demonstrated credentialed experience in IA therapy with adequate outcomes. It also is anticipated that after 2012, additional manpower for providing IA stroke therapy in stroke centers will come exclusively from fellowship-trained neurointerventionalists.

Training in Endovascular Surgical Neuroradiology

In the United States, Canada, Europe, and Japan, educational programs are specifically designed to train neuroscience physicians to treat acute hemorrhagic and ischemic stroke. This training represents the “gold standard” for performance of endovascular cerebrovascular procedures including the endovascular treatment of acute stroke.

Since the year 2000, program requirements have been formally established and published in North America to ensure uniform training in endovascular surgical neuroradiology, a field specializing in the endovascular treatment of acute stroke.¹⁷ “Endovascular surgical neuroradiology”

is the title or moniker used to describe the training pathway recognized by the ACGME.^{17,18}

Mechanical revascularization (ie, thrombectomy/embolectomy) devices have not yet been proven to improve patient outcomes. In addition, it is not possible to define general training requirements when there currently is significant clinical experience with only one mechanical revascularization clot-retrieval device and the necessary interventional skills for newer generations of clot-retrieval devices are unknown. Although there is consensus that the required skills and associated risks are greater for these devices than for catheter-based pharmacologic lysis, it is not possible to make specific training recommendations at this time. Nonetheless, at a minimum, the physician must meet the training criteria described herein for pharmacologic lysis for emergency stroke therapy and have successfully completed a training course for use of any specific device. Furthermore, the rates of procedural complications, including intracerebral hemorrhage, should conform to evidence-based national guidelines.

Maintenance of Physician and Facility Qualifications

Physicians should have ongoing stroke-specific continuing medical education (CME) of at least 15 hours every 2 years. Physicians also should have procedural outcomes that conform to national standards and institutional requirements.

Neuroendovascular procedures for acute ischemic stroke should be performed only at Joint Commission- or state-certified primary or comprehensive stroke centers.¹⁹ Outlying and community hospitals should develop access via telemedicine or other means to obtain endovascular acute stroke therapy through the development of stroke systems of care if comprehensive stroke center capabilities are not available.²⁰ Interventionalists must have 24/7 access to neurologists and neurosurgeons knowledgeable in patient selection and in the preneurologic and postneurologic critical care of endovascularly treated stroke patients. There must be 24/7 neurology and neurosurgical availability to treat possible complications of stroke therapy. There must be an active quality assurance program for stroke therapy to monitor outcomes both in the periprocedural period and at 90 days. All patients who have received emergency intervention for stroke must be reviewed by this quality assurance program. Outcomes should be tracked and recorded. All centers should participate in and enroll all stroke patients in available national stroke registries or multicenter stroke trials.

Conclusions

Medical specialties with ACGME-approved training in neuroscience and the care of stroke patients agree on the

importance of safety and quality of care for stroke patients. Due to the grave consequences of inadequate or deficient training, stringent credentialing criteria with formal neuroscience training as specified herein and by peer-reviewed published standards should be mandated for all providers of emergency endovascular stroke therapy,^{17,21} analogous to vascular interventions for acute myocardial infarction or other conditions with high morbidity and mortality.²²⁻³⁰ Credentialing committees at each health care facility are empowered to enforce training and practice standards and thus have an obligation to maintain recognized accreditation standards and to be aware of recommendations endorsed by the national organizations most directly involved in the diagnosis and management of acute stroke. Physician credentials, maintenance of certification, and quality improvement programs must be consistent with mandated and accepted standards defined by the ACGME, American Medical Association (AMA), ABMS, and state licensing boards.

References

- Rosamond W, Flegal K, Furie K, et al. Heart disease and stroke statistics—2008 update: A report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 2008;117:e25-e146.
- Sacco RL, Benjamin EJ, Broderick JP, et al. American Heart Association Prevention Conference, IV. Prevention and rehabilitation of stroke: risk factors. *Stroke* 1997; 28:1507-1517.
- Adelman SM. The National Survey of Stroke: economic impact. *Stroke* 1981;12:169-187.
- Taylor TN, Davis PH, Torner JC, et al. Lifetime cost of stroke in the United States. *Stroke* 1996;27:1459-1466.
- National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. Tissue plasminogen activator for acute ischemic stroke. *N Engl J Med* 1995;333: 1581-1587.
- del Zoppo GJ, Higashida RT, Furlan AJ, et al. PROACT: A phase II randomized trial of recombinant prourokinase by direct arterial delivery in acute middle cerebral artery stroke. *Stroke* 1998;29:4-11.
- Furlan A, Higashida R, Wechsler L, et al. Intra-arterial prourokinase for acute ischemic stroke. The PROACT II Study: a randomized controlled trial. *JAMA* 1999; 282:2003-2011.
- Smith WS, Sung G, Starkman S, et al. Safety and efficacy of mechanical embolectomy in acute ischemic stroke: results of the Merci Trial. *Stroke* 2005;36:1432-1438.
- Mas JL, Chatellier G, Beyssen B, et al. Endarterectomy versus stenting in patients with symptomatic severe carotid stenosis. *N Engl J Med* 2006;355:1660-1671.
- Ringleb PA, Allenberg J, Bruckmann H, et al. 30-day results from the space trial of stent-protected angioplasty versus carotid endarterectomy in symptomatic patients: a randomised non-inferiority trial. *Lancet* 2006;368: 1239-1247.
- American College of Cardiology (ACC). Practice guideline for the performance of diagnostic cervicocerebral angiography in adults. Amended 2007; (Resolution 12m:15) Available from: <http://www.acr.org>
- Adams HP Jr, del Zoppo G, Alberts MJ, et al. Guidelines for the early management of adults with ischemic stroke: a guideline from the American Heart Association/American Stroke Association Stroke Council, Clinical Cardiology Council, Cardiovascular Radiology and Intervention Council, and the Atherosclerotic Peripheral Vascular Disease and Quality of Care Outcomes in Research Interdisciplinary Working Groups. *Circulation* 2007; 115:e478-e534.
- Connors JJ 3rd, Sacks D, Furlan AJ, et al. Training, competency, and credentialing standards for diagnostic cervicocerebral angiography, carotid stenting, and cerebrovascular intervention. A joint statement from the American Academy of Neurology, American Association of Neurological Surgeons Cerebrovascular Section, American Society of Interventional and Therapeutic Neuroradiology, American Society of Neuroradiology, Congress of Neurological Surgeons, and Society of Interventional Radiology. *J Vasc Interv Radiol* 2004; 15:1347-1356.
- Spies JB, Sacks D. Credentials for uterine artery embolization. *J Vasc Interv Radiol* 2004;15:111-113.
- King SB 3rd, Aversano T, Ballard WL, et al. ACCF/AHA/SCAI 2007 update of the clinical competence statement on cardiac interventional procedures. A report of the American College of Cardiology Foundation/American Heart Association/American College of Physicians Task Force on Clinical Competence and Training. *J Am Coll Cardiol* 2007;50:82-108.
- Quality improvement guidelines for adult diagnostic neuroangiography. Cooperative study between the ASNR, ASITN, and SCVIR. *AJNR Am J Neuroradiol* 2000;21:146-150.
- Higashida RT, Hopkins LN, Berenstein A, et al. Program requirements for residency/fellowship education in neuroendovascular surgery/interventional neuroradiology: a special report on graduate medical education. *AJNR Am J Neuroradiol* 2000;21:1153-1159.
- American Academy of Neurology (AAN). Stroke and vascular neurology fellowship core curriculum. *Stroke* Section. 2001:1-5. Available from: <http://www.aan.com>.
- Adams R, Acker J, Alberts M, et al. Recommendations for improving the quality of care through stroke centers and systems: an examination of stroke center identification options. *Stroke* 2002;33:e1-e7.
- Schwamm LH, Holloway RG, Amarenco P, et al. Review of the evidence for the use of telemedicine within stroke systems of care. *Stroke* 2009;40:2635-2660.
- Barr JD, Connors JJ 3rd, Sacks D, et al. Quality improvement guidelines for the performance of cervical carotid angioplasty and stent placement. *AJNR Am J Neuroradiol* 2003;24:2020-2034.
- Pepine CJ, Allen HD, Bashore TM, et al. ACC/AHA guidelines for cardiac catheterization and cardiac catheterization laboratories. American College of Cardiology/American Heart Association Ad Hoc Task Force on Cardiac Catheterization. *Circulation* 1991; 84:2213-2247.
- Friesinger GC, Adams DF, Bourassa MG, et al. Optimal resources for examination of the heart and lungs: cardiac catheterization and radiographic facilities. Examination of the Chest and Cardiovascular System Study Group. *Circulation* 1983;68:891A-930A.
- Conti CR, Faxon DP, Gruentzig A, et al. 17th Bethesda Conference: adult cardiology training. Task force III: training in cardiac catheterization. *J Am Coll Cardiol* 1986;7:1205-1206.

25. Hodgson JM, Tommaso CL, Watson RM, et al. Core curriculum for the training of adult invasive cardiologists. Report of the Society for Cardiac Angiography and Interventions Committee on Training Standards. *Cathet Cardiovasc Diagn* 1996;37:392-408.
26. Hirshfeld JW Jr, Ellis SG, Faxon DP. Recommendations for the assessment and maintenance of proficiency in coronary interventional procedures. Statement of the American College of Cardiology. *J Am Coll Cardiol* 1998; 31:722-743.
27. Ryan TJ, Faxon DP, Gunnar RM, et al. Guidelines for percutaneous transluminal coronary angioplasty. A report of the American College of Cardiology/American Heart Association Task Force on Assessment of Diagnostic and Therapeutic Cardiovascular Procedures (Subcommittee on Percutaneous Transluminal Coronary Angioplasty). *Circulation* 1988;78:486-502.
28. Ryan TJ, Klocke FJ, Reynolds WA. Clinical competence in percutaneous transluminal coronary angioplasty. A statement for physicians from the ACP/ACC/AHA Task Force on Clinical Privileges in Cardiology. *J Am Coll Cardiol* 1990;15:1469-1474.
29. Ryan TJ, Bauman WB, Kennedy JW, et al. Guidelines for percutaneous transluminal coronary angioplasty. A report of the American Heart Association/American College of Cardiology Task Force on Assessment of Diagnostic and Therapeutic Cardiovascular Procedures (Committee on Percutaneous Transluminal Coronary Angioplasty). *Circulation* 1993;88:2987-3007.
30. Cowley MJ, Faxon DP, Holmes DR Jr. Guidelines for training, credentialing, and maintenance of competence for the performance of coronary angioplasty. A report from the Interventional Cardiology Committee and the Training Program Standards Committee of the Society for Cardiac Angiography and Interventions. *Cathet Cardiovasc Diagn* 1993;30:1-4.